



Pre-Admission Labs and Diagnostics

Pre-Surgery Diagnosis: _____

Scheduled Procedure: _____

Date of Procedure: _____ Allergies: _____

Precertification #: _____

- No Lab Tests Required
- Anesthesiology Consultation (Patients with Medical or Surgical issues) MRSA Screen
- Hgb/Hct Urine HCG Type and Screen
- CBC without diff Serum HCG Crossmatch ___# units PRBCs
- CBC w/diff Quant. HCG Crossmatch ___# units autologous Blood
- PT HBsAg
- PTT Sickle Cell Screening Other blood products _____
- CMP Amylase CXR
- BMP Lipase KUB
- Blood Glucose Hepatic Panel Other X-ray _____
- FSBS Day of Surgery Other Lab Ultrasound _____
- Uric Acid PFT
- Urinalysis w/ Micro

Blood for pre-transfusion testing must be drawn within 48 hours of surgery. Typed and Screened blood can only be held 48 hours.

Pre-Surgery Orders OPS (Outpatient Surgery) IP (Inpatient) Unknown Length of Stay

Vital Signs: Per Protocol Other _____

Diet: NPO NPO after _____

- LR at KVO or 100ml/hr 200ml/hr _____ml/hr
- NS at KVO or 100ml/hr 200ml/hr _____ml/hr
- NS (500ml bag) at KVO via microdrip tubing for renal patient

Medications:

- Pre-Op Antibiotic in ASU/OR holding area _____
- Bacterial Endocarditis Prophylaxis: _____ Consult Pharmacy/Anesthesia
- Ancef 1GM 2GM IV in OR
- If PCN / B-Lactam Allergy OR Patient at Increased risk for infection
- Vancomycin 1GM IV to be started within 120 minutes of incision or
- Cleocin 900MG IV to be started within 30 minutes of incision
- Tylenol suppository by weight in ASU/OR
- Other _____

Consult Anesthesia for post-op pain block
(Please note below if you would like a specific type of block considered for this patient)

Preparation:

- Incentive Spirometry: Instructions/pre-admit
- Thigh / Knee High TED hose
- Sequential Compression Device in OR

Minimum Testing Guidelines (Anesthesia Service): These guidelines are suggested minimums. They are not replacement for medical judgment, and the patient's medical history and/or the proposed surgical procedure may indicate additional laboratory or diagnostic testing.

No pre-operative laboratory testing is required for asymptomatic patients without significant medical problems who are less than 40 years of age.

ECG: Males aged 40 and above require an ECG. Females aged 50 and above require ECG

Pregnancy Test: A pregnancy test is required for all menstruating females, unless not indicated due to sterility. Serum HCG if blood is drawn for other tests, otherwise it will be a urine HCG.

Labs: CBC and BMP are required for patients age 65 and older. BMP is required for diabetic, renal and/or hypertensive patients. FSBS is required day of surgery in diabetic patients, except in patient who have a BMP done day of surgery.

Copies: A copy of a CXR and/or ECG completed in the past 6 months is sufficient in the absence of a change in the patient's health status. A copy of lab work completed in the past 30 days is sufficient in the absence of renal disease. Renal patients must have a K+ completed after their last dialysis treatment before the date of surgery.

IVF: Every patient over age 10 is required to have an IV of LR at KVO rate. Renal patients must have NS at KVO rate