



ROUTINE PRE-OP TAH / LAVH ORDERS

Pre-Admission Labs and Diagnostics

Pre-Surgery Diagnosis: _____

Scheduled Procedure: _____

Date of Procedure: _____ Allergies: _____

Precertification #: _____

- No Lab Tests Required
- Anesthesiology Consultation(Patients with Medical or Surgical issues)
- Hgb/Hct
- CBC w/diff
- PT
- PTT
- Blood Glucose
- Electrolytes-Na,K+,Cl,CO2
- BMP
- Urinalysis
- Pregnancy Test
- HBsAg
- Sickle Cell Screening
- Amylase
- Hepatic Panel
- ECG
- CXR
- X-ray Type _____
- Ultrasound Type _____
- Type & Screen
- T&S (Crossmatch) ___# units PRBCs
- T&S (Crossmatch) ___# units autologous Blood

Blood for pre-transfusion testing must be drawn within 48 hours of surgery. Typed and Screened blood can only be held 48 hours.

Pre-Surgery Orders OPS (Outpatient Surgery) IP (Inpatient)

Vital Signs: Per Protocol Other _____

Diet: NPO NPO after _____ LR at _____ 100ml/hr 200ml/hr ___ml/hr

Medications:

- Pre-Op Antibiotic: _____
- Bacterial Endocarditis Prophylaxis: _____ Consult Pharmacy/Anesthesia
- Ancef 1GM 2GM IV in OR
- Unasyn 1.5GM 3GM IV in OR
- If PCN / B-Lactam Allergy:
- Cleocin 900MG + Gentamycin 80MG IV 60 minutes prior to incision.
- Bicitra 30ml po on-call Other _____

Preparation:

- Incentive Spirometry Instructions/pre-admit
- Thigh / Knee High TED hose
- Sequential Compression Device in OR
- Enema SS on Admit
- Betadine Douche pre-op

Miscellaneous:

- H&P
- To be done on admit by _____
- Consult obtained from _____

Minimum Testing Guidelines (Anesthesia Service): These guidelines are suggested minimums.

They are not replacement for medical judgment, and the patient's medical history and/or the proposed surgical procedure may indicate additional laboratory or diagnostic testing.

No pre-operative laboratory testing is required for asymptomatic patients without significant medical problems who are less than 40 years of age, except for an Hgb/Hct for 0-6 months as per protocol, and except for urine HCG.

ECG: Males, aged 40 and above require an ECG. Males and Females aged 50 and above require ECG

CXR: CXRs are not required in the absence of cardiorespiratory disease. Inpatients with cardiorespiratory disease, a CXR with the past six months is sufficient in the absence of a significant change in status of the cardiorespiratory illness.

Pregnancy Test: A pregnancy test is required for all menstruating females scheduled for anesthesia or surgery, unless not indicated due to sterility. This test will be a serum HCG if blood is drawn for other tests. Otherwise it will be a urine HCG.

Patient Name and DOB

Physician Offices—
For the most updated form please visit
www.coffeeregional.org and
print from the "For Our Physicians" link.